arXiv:1505.01590v1 [math.GT] 7 May 2015

GENERALIZED TORSION ELEMENTS IN THE KNOT GROUPS OF TWIST KNOTS

MASAKAZU TERAGAITO

Abstract. It is well known that any knot group is torsion-free, but it may admit a generalized torsion element. We show that the knot group of any negative twist knot admits a generalized torsion element. This is a generalization of the same claim for the knot 5_2, which is the (-2)-twist knot, by Naylor and Rolfsen.

1. Introduction

For a knot K in the 3-sphere S^3, the *knot group* of K is the fundamental group of the complement $S^3 - K$. It is a classical fact that any knot group is torsion-free. However, many knot groups can have generalized torsion elements. For a group, a non-trivial element is called a *generalized torsion element* if some non-empty finite product of its conjugates is the identity. Typical exmples are the knot groups of torus knots (see [6]). It was open whether the knot group of a hyperbolic knot admits a generalized torsion element or not. However, Naylor and Rolfsen [6] first found such an element in the knot group of the hyperbolic knot 5_2. As they wrote, the element was found with the help of computer, so a (topological) meaning of the element is not obvious.

The knot 5_2 is the (-2)-twist knot. See below for the convention of twist knots. In this article, we completely determine the existence of generalized torsion elements in the knot groups of twist knots as a generalization of Naylor-Rolfsen's result. Figure 1 shows the m-twist knot, where the rectangle box contains the right-handed (resp. left-handed) horizontal m-full twists if $m > 0$ (resp. $m < 0$). By this convention, the 1-twist knot is the figure-eight knot, and the (-1)-twist knot is the right-handed trefoil as shown in Figure 1. We may call the m-twist knot a positive or negative twist knot, according to the sign of m.



Figure 1. The m-twist knot and the (-1)-twist knot

2010 *Mathematics Subject Classification.* Primary 57M25; Secondary 57M05, 06F15.

The author was partially supported by Japan Society for the Promotion of Science, Grant-in-Aid for Scientific Research (C), 25400093.



Figure 2. A Seifert surface ($n = 2$)

There is a well known relation between the bi-orderability and the existence of generalized torsion elements in a group. We recall that a group is said to be *bi-orderable* if it admits a strict total ordering, which is invariant under multiplication on both sides. Then any bi-orderable group has no generalized torsion elements ([6]). The converse statement is not true, in general ([5]).

According to [2], the knot group of any positive twist knot is bi-orderable, but that of any negative twist knot is not bi-orderable. Therefore, the former does not admit a generalized torsion element, but we can expect that the latter would admit it. In fact, Naylor and Rolfsen [6] confirmed this claim for the (-2)-twist knot. The main result of the present article is to verify this expectation.

Theorem 1.1. *The knot group of any negative twist knot admits a generalized torsion element.*

Throughout the paper, we use the convention $x^g = g^{-1}xg$ for the conjugate of x by g and $[x, y] = x^{-1}y^{-1}xy$ for the commutator.

2. Presentations of knot groups

For $n \geq 1$, let K be the $(-n)$-twist knot. We prepare a certain presentation of the knot group of K by using its Seifert surface. Figure 2 shows a Seifert surface S where $n = 2$. The fundamental group $\pi_1(S)$ is a free group of rank two, generated by x and y as illustrated there. Since the complement of the regular neighborhood $N(S)$ of S in S^3 is a genus two handlebody, $\pi_1(S^3 - N(S))$ is also a free group of rank two, generated by a and b.

Theorem 2.1. *The knot group G of the $(-n)$-twist knot K has a presentation*

$$\langle a, b, t \mid tat^{-1} = b^{-1}a, t(b^n a^{-1})t^{-1} = b^n \rangle,$$

where t is a meridian.

Proof. If we push x and y off from the Seifert surface to the front side, then we obtain $x^+ = a$ and $y^+ = b^n a^{-1}$. Similarly, if we push them off to the back side, then $x^- = b^{-1}a$ and $y^- = b^n$. As shown in [4, Lemma 2.1] (it is just an application of the van Kampen theorem), the knot group G is generated by a, b and a meridian t with two relations $tx^+t^{-1} = x^-$ and $ty^+t^{-1} = y^-$. □

From the second relation, we have

$$a = [t, b^n]. \tag{2.1}$$

Thus the knot group G is generated by b and t.

Remark 2.2. This presentation is distinct from one used in [6] when $n = 2$, but it is equivalent, of course.

By their choices, the generators a and b belong to the commutator subgroup of G. Furthermore, we have $a = [t, b^n]$ and $b = [a^{-1}, t^{-1}]$. This implies that these elements belong to the intersection of all members of the lower central series of G.

Lemma 2.3. *Let $D = [t, b]$. Then $D \neq 1$.*

Proof. If $D = 1$, then the knot group G would be abelian, because G is generated by t and b. Since the knot is non-trivial, this is impossible. □

Lemma 2.4. $a = DD^b \cdots D^{b^{n-1}}$.

Proof. In general, $[x, y^m] = [x, y][x, y^{m-1}]^y$. The conclusion immediately follows from this and (2.1). □

3. Proof of Theorem 1.1

In this section, we prove that $D = [t, b]$ is a generalized torsion element of G. Let $\langle D \rangle$ be the semigroup consisting of all non-empty products of conjugates of D in G. That is,

$$\langle D \rangle = \{D^{g_1} D^{g_2} \cdots D^{g_m} \mid g_1, g_2, \dots, g_m \in G, m \geq 1\}.$$

If the semigroup $\langle D \rangle$ is shown to contain the identity, then we can conclude that D is a generalized torsion element.

From the second relation of the presentation in Theorem 2.1, $(tb^n t^{-1})(ta^{-1}t^{-1}) = b^n$. Then $tb^n t^{-1} = b^n(tat^{-1})$. By the first relation of the presentation, we have $tb^n t^{-1} = b^{n-1}a$. Thus, by (2.1), we have

$$tb^n t^{-1} = b^{n-1} t^{-1} b^{-n} t b^n. \tag{3.1}$$

This can also be modified to

$$b^{-(n-1)} t b^n t^{-1} = [t, b^n]. \tag{3.2}$$

Proposition 3.1. *$b^n \in \langle D \rangle$.*

Proof. We show $tb^n t^{-1} \in \langle D \rangle$, which imples $b^n \in \langle D \rangle$. By (3.1), it is sufficient to show $b^{n-1} t^{-1} b^{-n} t b^n \in \langle D \rangle$.

Conjugate $b^{n-1} t^{-1} b^{-n} t b^n$ by $t^{-(n-1)}$. By applying (3.2) repeatedly, we have

$$\begin{aligned}
t^{n-1}(b^{n-1}t^{-1}b^{-n}tb^n)t^{-(n-1)} &= t^{n-1}b^{n-1}(t^{-1}b^{-1}) \cdot b^{-(n-1)}tb^n t^{-1} \cdot t^{-(n-2)} \\
&= t^{n-1}b^{n-1}(t^{-1}b^{-1}) \cdot [t, b^n] \cdot t^{-(n-2)} \\
&= t^{n-1}b^{n-1}(t^{-1}b^{-1})^2 \cdot b^{-(n-1)}tb^n t^{-1} \cdot t^{-(n-3)} \\
&= t^{n-1}b^{n-1}(t^{-1}b^{-1})^2 \cdot [t, b^n] \cdot t^{-(n-3)} \\
&\ \ \vdots \\
&= t^{n-1}b^{n-1}(t^{-1}b^{-1})^{n-1} \cdot [t, b^n].
\end{aligned}$$

Since $[t, b^n] = a \in \langle D \rangle$, we can stop here when $n = 1$. Hereafter, we suppose $n \geq 2$. Then, it suffices to show $t^{n-1}b^{n-1}(t^{-1}b^{-1})^{n-1} \in \langle D \rangle$.

Claim 3.2. $t^{n-1}b^{n-1}(t^{-1}b^{-1})^{n-1} = D^{(bt)^{n-2}b^{-(n-1)}t^{-(n-1)}} \cdot t^{n-1}b^{n-1}(t^{-1}b^{-1})^{n-2}b^{-1}t^{-1}$.

Proof of Claim 3.2. This follows from

$$\begin{aligned} D^{(bt)^{n-2}b^{-(n-1)}t^{-(n-1)}} &= t^{n-1}b^{n-1}(t^{-1}b^{-1})^{n-2}D(bt)^{n-2}b^{-(n-1)}t^{-(n-1)} \\ &= t^{n-1}b^{n-1}(t^{-1}b^{-1})^{n-2}\cdot t^{-1}b^{-1}tb\cdot (bt)^{n-2}b^{-(n-1)}t^{-(n-1)} \\ &= t^{n-1}b^{n-1}(t^{-1}b^{-1})^{n-1}\cdot tb(bt)^{n-2}b^{-(n-1)}t^{-(n-1)}. \end{aligned}$$

□

If $n=2$, then the right hand side is $D^{b^{-1}t^{-1}}$, so we are done.

Claim 3.3. *If $n\geq 3$, then*

$$t^{n-1}b^{n-1}(t^{-1}b^{-1})^{n-2}b^{-1}t^{-1} = [t^{-1},b^{-(n-1)}]^{t^{-(n-2)}}\cdots[t^{-1},b^{-3}]^{t^{-2}}[t^{-1},b^{-2}]^{t^{-1}}.$$

Proof of Claim 3.3. We use an induction on n. When $n=3$, the left hand side is $t^2b^2(t^{-1}b^{-1})b^{-1}t^{-1}$, which is equal to $[t^{-1},b^{-2}]^{t^{-1}}$.

Assume that

$$t^{n-2}b^{n-2}(t^{-1}b^{-1})^{n-3}b^{-1}t^{-1} = [t^{-1},b^{-(n-2)}]^{t^{-(n-3)}}\cdots[t^{-1},b^{-3}]^{t^{-2}}[t^{-1},b^{-2}]^{t^{-1}}.$$

Multiplying the left hand side $[t^{-1},b^{-(n-1)}]^{t^{-(n-2)}}$ from the left gives

$$[t^{-1},b^{-(n-1)}]^{t^{-(n-2)}}t^{n-2}b^{n-2}(t^{-1}b^{-1})^{n-3}b^{-1}t^{-1} = t^{n-1}b^{n-1}(t^{-1}b^{-1})^{n-2}b^{-1}t^{-1}.$$

□

Recall that $[t,b^m]\in\langle D\rangle$ for $m\geq 1$, as in the proof of Lemma 2.4. Also, $[t^{-1},b^{-m}]=[t,b^m]^{t^{-1}b^{-m}}\in\langle D\rangle$. By Claims 3.2 and 3.3, $t^{n-1}b^{n-1}(t^{-1}b^{-1})^{n-1}\in\langle D\rangle$. This completes the proof. □

Proposition 3.4. $b^{-n}\in\langle D\rangle$.

Proof. The argument is similar to that of the proof of Proposition 3.1. The relation (3.1) can be changed into

$$t^{-1}b^{-n}t = b^{-(n-1)}tb^nt^{-1}b^{-n}. \tag{3.3}$$

We can see that the relation (3.1) is invariant under the transformation $t\to t^{-1}$, $b\to b^{-1}$. Thus, conjugating (3.3) by t^{n-1} yields the relation

$$t^{-n}b^{-n}t^n = t^{-(n-1)}b^{-(n-1)}(tb)^{n-1}\cdot[t^{-1},b^{-n}].$$

When $n=1$, the right hand side is $[t^{-1},b^{-1}]=D^{b^{-1}t^{-1}}\in\langle D\rangle$. So, assume $n\geq 2$. As above, $[t^{-1},b^{-n}]\in\langle D\rangle$.

Claim 3.5. $t^{-(n-1)}b^{-(n-1)}(tb)^{n-1} = D^{(b^{-1}t^{-1})^{n-1}b^{n-1}t^{n-1}}\cdot t^{-(n-1)}b^{-(n-1)}(tb)^{n-2}bt.$

Proof of Claim 3.5. By the transformation $t\to t^{-1}$, $b\to b^{-1}$, the relation of Claim 3.2 changes into

$$t^{-(n-1)}b^{-(n-1)}(tb)^{n-1} = [t^{-1},b^{-1}]^{(b^{-1}t^{-1})^{n-2}b^{n-1}t^{n-1}}\cdot t^{-(n-1)}b^{-(n-1)}(tb)^{n-2}bt.$$

Since $[t^{-1},b^{-1}]=D^{b^{-1}t^{-1}}$, we have the conclusion. □

Again, if $n=2$, then the right hand side is $D^{b^{-1}t^{-1}bt}=D$.

Claim 3.6. *If $n\geq 3$, then*

$$t^{-(n-1)}b^{-(n-1)}(tb)^{n-2}bt = [t,b^{n-1}]^{t^{n-2}}\cdots[t,b^3]^{t^2}[t,b^2]^t.$$

Proof. This can be proved on an induction on n as in the proof of Claim 3.3. We omit this. ☐

Since $[t, b^m] \in \langle D\rangle$ for $m \geq 1$, we obtain $b^{-n} \in \langle D\rangle$. ☐

Proof of Theorem 1.1. By Propositions 3.1 and 3.4, both of b^n and b^{-n} belong to the semigroup $\langle D\rangle$. Hence $\langle D\rangle$ contains the identity. Since $D \neq 1$ by Lemma 2.3, D is a generalized torsion element. ☐

Remark 3.7. For the case $n = 2$, Naylor and Rolfsen [6] showed that $[b^{-1}, t]$ in our notation is a generalized torsion element. This equals to $D^{b^{-1}}$, which is a conjugate of our D.

4. Comments

A group without generalized torsion elements is called an R^*-group in literature ([5]). It was an open question whether the class of bi-orderable groups coincided with the class of R^*-groups [5, p.79]. Unfortunately, the answer is known to be negative. However, there might be a possibility that two classes coincide among knot groups. Thus we expect that any non-bi-orderable knot group would admit a generalized torsion element. Many knot groups are now known to be non-bi-orderable by [1, 2, 3, 7]. It would be a next problem to examine such knot groups.

Acknowledgments

The author would like to thank Makoto Sakuma for helpful conversations.

References

1. I. Chiswell, A. Glass and J. Wilson, *Residual nilpotence and ordering in one-relator groups and knot groups*, preprint.
2. A. Clay, C. Desmarais and P. Naylor, *Testing bi-orderability of knot groups*, preprint.
3. A. Clay and D. Rolfsen, *Ordered groups, eigenvalues, knots, surgery and L-spaces*, Math. Proc. Cambridge Philos. Soc. **152** (2012), no. 1, 115–129.
4. X. S. Lin, *Representations of knot groups and twisted Alexander polynomials*, Acta Math. Sin. **17** (2001), no. 3, 361–380.
5. R. Mura and A. Rhemtulla, *Orderable groups*, Lecture Notes in Pure and Applied Mathematics, Vol. 27. Marcel Dekker, Inc., New York-Basel, 1977.
6. G. Naylor and D. Rolfsen, *Generalized torsion in knot groups*, preprint.
7. B. Perron and D. Rolfsen, *On orderability of fibred knot groups*, Math. Proc. Camb. Phil. Soc. **135** (2003), 147–153.

Department of Mathematics and Mathematics Education, Hiroshima University, 1-1-1 Kagamiyama, Higashi-hiroshima, Japan 739-8524.
E-mail address: `teragai@hiroshima-u.ac.jp`